arXiv:1509.09193v1 [math.NT] 30 Sep 2015

IDENTITIES OF SYMMETRY FOR THE GENERALIZED DEGENERATE EULER POLYNOMIALS

DAE SAN KIM AND TAEKYUN KIM

Abstract. In this paper, we give some identities of symmetry for the generalized degenerate Euler polynomials attached to χ which are derived from the symmetric properties for certain fermionic p-adic integrals on $\mathbb{Z}_p$.

1. Introduction and preliminaries

Let p be a fixed odd prime. Throughout this paper, $\mathbb{Z}_p, \mathbb{Q}_p$ and $\mathbb{C}_p$ will be the ring of p-adic integers, the field of p-adic rational numbers and the completion of the algebraic closure of $\mathbb{Q}_p$, respectively.

The p-adic norm $|\cdot|_p$ in $\mathbb{C}_p$ is normalized as $|p|_p = \frac{1}{p}$. Let $f(x)$ be continuous function on $\mathbb{Z}_p$. Then the fermionic p-adic integral on $\mathbb{Z}_p$ is defined as

$$I_{-1}(f) = \int_{\mathbb{Z}_p} f(x)\, d\mu_{-1}(x) = \lim_{N\to\infty} \sum_{x=0}^{p^N-1} f(x)(-1)^x, \quad (\text{see } [9]). \tag{1.1}$$

From (1.1), we note that

$$I_{-1}(f_n) + (-1)^{n-1} I_{-1}(f) = 2\sum_{l=0}^{n-1} (-1)^{n-1-l} f(l), \quad (\text{see } [7]), \tag{1.2}$$

where $n \in \mathbb{N}$.

As is well known, the Euler polynomials are defined by the generating function

$$\int_{\mathbb{Z}_p} e^{(x+y)t} d\mu_{-1}(y) = \frac{2}{e^t+1} e^{xt} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}. \tag{1.3}$$

When $x = 0$, $E_n = E_n(0)$ are called the Euler numbers (see [1–19]).

For a fixed odd integer d with $(p, d) = 1$, we set

$$X = \varprojlim_N \mathbb{Z}/dp^N\mathbb{Z}, \quad X^* = \bigcup_{\substack{0<a<dp \\ (a,p)=1}} (a + dp\mathbb{Z}_p),$$

$$a + dp^N\mathbb{Z}_p = \left\{ x \in X \mid x \equiv a \pmod{dp^N} \right\},$$

where $a \in \mathbb{Z}$ lies in $0 \le a < dp^N$.

2010 *Mathematics Subject Classification.* 11B68, 11B83, 11C08, 65D20, 65Q30, 65R20.

Key words and phrases. Identities of symmetry, Generalized degenerate Euler polynomial, fermionic p-adic integral.

It is known that

$$\int_{\mathbb{Z}_p} f(x)\, d\mu_{-1}(x) = \int_X f(x)\, d\mu_{-1}(x), \quad (\text{see } [7\text{–}9]),$$

where f is a continuous function on $\mathbb{Z}_p$.

Let $d \in \mathbb{N}$ with $d \equiv 1 \pmod 2$ and let χ be a Dirichlet character with conductor d. Then the generalized Euler polynomials attached to χ are defined by the generating function

$$\left(\frac{2\sum_{a=0}^{d-1}(-1)^a \chi(a)\, e^{at}}{e^{dt}+1}\right) e^{xt} = \sum_{n=0}^{\infty} E_{n,\chi}(x)\frac{t^n}{n!}. \tag{1.4}$$

In particular, for $x = 0$, $E_{n,\chi} = E_{n,\chi}(0)$ are called the generalized Euler numbers attached to χ.

For $d \in \mathbb{N}$ with $d \equiv 1 \pmod 2$, by (1.2), we get

$$\begin{aligned} &\int_X \chi(y)\, e^{(x+y)t} d\mu_{-1}(y) \\ &= \frac{2\sum_{a=0}^{d-1}(-1)^a \chi(a)\, e^{at}}{e^{dt}+1} e^{xt} \\ &= \sum_{n=0}^{\infty} E_{n,\chi}(x)\frac{t^n}{n!}, \quad (\text{see } [9\text{–}11]). \end{aligned} \tag{1.5}$$

From (1.5), we have

$$\int_X \chi(y)(x+y)^n\, d\mu_{-1}(y) = E_{n,\chi}(x), \quad (n \geq 0). \tag{1.6}$$

Carlitz considered the degenerate Euler polynomials given by the generating function

$$\begin{aligned} &\frac{2}{(1+\lambda t)^{\frac{1}{\lambda}}+1}(1+\lambda t)^{\frac{x}{\lambda}} \\ &= \sum_{n=0}^{\infty} \mathcal{E}_n(x \mid \lambda)\frac{t^n}{n!}, \quad (\text{see } [3]). \end{aligned} \tag{1.7}$$

Note that $\lim_{\lambda\to 0} \mathcal{E}_n(x \mid \lambda) = E_n(x)$, $(n \geq 0)$.

From (1.2), we note that

$$\begin{aligned} &\int_X (1+\lambda t)^{\frac{x+y}{\lambda}}\, d\mu_{-1}(y) \\ &= \frac{2}{(1+\lambda t)^{\frac{1}{\lambda}}+1}(1+\lambda t)^{\frac{x}{\lambda}} \\ &= \sum_{n=0}^{\infty} \mathcal{E}_n(x \mid \lambda)\frac{t^n}{n!}. \end{aligned} \tag{1.8}$$

Thus, by (1.8), we get

$$\int_X (y+x \mid \lambda)_n\, d\mu_{-1}(y) = \mathcal{E}_n(x \mid \lambda), \quad (n \geq 0), \tag{1.9}$$

where $(x \mid \lambda)_n = x(x-\lambda)\cdots(x-(n-1)\lambda)$, for $n \geq 1$, and $(x \mid \lambda)_0 = 1$.

From (1.2), we can derive the following equation:

$$\int_X \chi(y)(1+\lambda t)^{\frac{x+y}{\lambda}} d\mu_{-1}(y) = \frac{2\sum_{a=0}^{d-1}(-1)^a \chi(a)(1+\lambda t)^{\frac{a}{\lambda}}}{(1+\lambda t)^{\frac{d}{\lambda}}+1}(1+\lambda t)^{\frac{x}{\lambda}}, \tag{1.10}$$

where $d \in \mathbb{N}$ with $d \equiv 1 \pmod 2$.

In view of (1.5), we define the generalized degenerate Euler polynomials attached to χ as follows:

$$\frac{2\sum_{a=0}^{d-1}(-1)^a \chi(a)(1+\lambda t)^{\frac{a}{\lambda}}}{(1+\lambda t)^{\frac{d}{\lambda}}+1}(1+\lambda t)^{\frac{x}{\lambda}} = \sum_{n=0}^{\infty} \mathcal{E}_{n,\lambda,\chi}(x)\frac{t^n}{n!}. \tag{1.11}$$

When $x=0$, $\mathcal{E}_{n,\lambda,\chi} = \mathcal{E}_{n,\lambda,\chi}(0)$ are called the generalized degenerate Euler numbers attached to χ.

Let n be an odd natural number. Then, by (1.2), we get

$$\int_X \chi(x)(1+\lambda t)^{\frac{nd+x}{\lambda}} d\mu_{-1}(x) + \int_X \chi(x)(1+\lambda t)^{\frac{x}{\lambda}} d\mu_{-1}(x) = 2\sum_{l=0}^{nd-1}(-1)^l \chi(l)(1+\lambda t)^{\frac{l}{\lambda}}. \tag{1.12}$$

Now, we set

$$R_k(n,\lambda \mid x) = 2\sum_{l=0}^{n}(-1)^l \chi(l)(l \mid \lambda)_k. \tag{1.13}$$

From (1.2) and (1.12), we have

$$\begin{aligned}\int_X (1+\lambda t)^{\frac{x+dn}{\lambda}} \chi(x) d\mu_{-1}(x) + \int_X \chi(x)(1+\lambda t)^{\frac{x}{\lambda}} d\mu_{-1}(x) &= \frac{2\int_X (1+\lambda t)^{\frac{x}{\lambda}} \chi(x) d\mu_{-1}(x)}{\int_X (1+\lambda t)^{\frac{ndx}{\lambda}} d\mu_{-1}(x)} \\ &= \sum_{k=0}^{\infty} R_k(nd-1,\lambda \mid \chi)\frac{t^k}{k!},\end{aligned} \tag{1.14}$$

where $n, d \in \mathbb{N}$ with $n \equiv 1 \pmod 2$, $d \equiv 1 \pmod 2$.

In this paper, we give some identities of symmetry for the generalized degenerate Euler polynomials attached to χ derived from the symmetric properties of certain fermionic p-adic integrals on $\mathbb{Z}_p$.

2. Identities of symmetry for the generalized degenerate Euler polynomials

Let w_1, w_2 be an odd natural numbers. Then we consider the following integral equation:

$$\frac{\int_X \int_X (1+\lambda t)^{\frac{w_1 x_1 + w_2 x_2}{\lambda}} \chi(x_1)\chi(x_2) d\mu_{-1}(x_1) d\mu_{-1}(x_2)}{\int_X (1+\lambda t)^{\frac{dw_1 w_2 x}{\lambda}} d\mu_{-1}(x)} \tag{2.1}$$

$$= \frac{2\left((1+\lambda t)^{\frac{dw_1w_2}{\lambda}}+1\right)}{\left((1+\lambda t)^{\frac{w_1d}{\lambda}}+1\right)\left((1+\lambda t)^{\frac{w_2d}{\lambda}}+1\right)} \times \sum_{a=0}^{d-1}\chi(a)(1+\lambda t)^{\frac{w_1a}{\lambda}}(-1)^a \times \sum_{b=0}^{d-1}\chi(b)(1+\lambda t)^{\frac{w_2b}{\lambda}}(-1)^b.$$

From (1.10) and (1.11), we note that

$$\int_X \chi(y)(x+y\mid\lambda)_n\, d\mu_{-1}(y) = \mathcal{E}_{n,\lambda,\chi}(x), \quad (n\geq 0). \tag{2.2}$$

By (1.14), we get

$$\int_X \chi(x)(x+dn\mid\lambda)_k\, d\mu_{-1}(x) + \int_X \chi(x)(x\mid\lambda)_k\, d\mu_{-1}(x) = R_k(nd-1,\lambda\mid x), \tag{2.3}$$

where $k\geq 0$.

Thus, by (2.2) and (2.3), we get

$$\mathcal{E}_{k,\lambda,\chi}(nd) + \mathcal{E}_{k,\lambda,\chi} = R_k(nd-1,\lambda\mid\chi), \tag{2.4}$$

where $k\geq 0$, $n,d\in\mathbb{N}$ with $n\equiv 1 \pmod 2$, $d\equiv 1 \pmod 2$.

Now, we set

$$I_\chi(w_1,w_2\mid\lambda) = \frac{\int_X\int_X \chi(x_1)\chi(x_2)(1+\lambda t)^{\frac{w_1x_1+w_2x_2+w_1w_2x}{\lambda}}\, d\mu_{-1}(x_1)\, d\mu_{-1}(x_2)}{\int_X (1+\lambda t)^{\frac{dw_1w_2x}{\lambda}}\, d\mu_{-1}(x)}. \tag{2.5}$$

From (2.5), we have

$$I_\chi(w_1,w_2\mid\lambda) = \frac{2\left((1+\lambda t)^{\frac{dw_1w_2}{\lambda}}+1\right)(1+\lambda t)^{\frac{w_1w_2x}{\lambda}}}{\left((1+\lambda t)^{\frac{w_1d}{\lambda}}+1\right)\left((1+\lambda t)^{\frac{w_2d}{\lambda}}+1\right)} \times \sum_{a=0}^{d-1}\chi(a)(-1)^a(1+\lambda t)^{\frac{w_1a}{\lambda}} \times \sum_{b=0}^{d-1}\chi(b)(-1)^b(1+\lambda t)^{\frac{w_2b}{\lambda}}. \tag{2.6}$$

Thus, by (2.6), we see that $I_\chi(w_1,w_2\mid\lambda)$ is symmetric in w_1,w_2. By (1.12), (1.14), (2.2) and (2.5), we get

$$2I_\chi(w_1,w_2\mid\lambda) = \sum_{l=0}^{\infty}\left(\sum_{i=0}^{l}\binom{l}{i}\mathcal{E}_{i,\frac{\lambda}{w_2},\chi}(w_1x)\, w_2^i w_1^{l-i} R\left(dw_2-1,\frac{\lambda}{w_1}\middle|\chi\right)\right)\frac{t^l}{l!}. \tag{2.7}$$

From the symmetric property of $I_\chi(w_1,w_2\mid\lambda)$ in w_1 and w_2, we have

$$2I_\chi(w_1,w_2\mid\lambda) \tag{2.8}$$

$$= 2I_{\chi}\left(w_2, w_1 \mid \chi\right)$$
$$= \sum_{l=0}^{\infty}\left(\sum_{i=0}^{l}\binom{l}{i}\mathcal{E}_{i,\frac{\lambda}{w_1},\chi}\left(w_2x\right)w_1^i w_2^{l-i}R\left(dw_1-1,\frac{\lambda}{w_2}\middle|\chi\right)\right)\frac{t^l}{l!}.$$

Therefore, by (2.7) and (2.8), we obtain the following theorem.

Theorem 1. *For $w_1, w_2, d \in \mathbb{N}$ with $w_1 \equiv w_2 \equiv d \equiv 1 \pmod 2$, let χ be a Dirichlet character with conductor d. Then, we have*

$$\sum_{i=0}^{l}\binom{l}{i}\mathcal{E}_{i,\frac{\lambda}{w_1},\chi}\left(w_2x\right)w_1^i w_2^{l-i}R\left(dw_1-1,\frac{\lambda}{w_2}\middle|\chi\right)$$
$$= \sum_{i=0}^{l}\binom{l}{i}\mathcal{E}_{i,\frac{\lambda}{w_2},\chi}\left(w_1x\right)w_2^i w_1^{l-i}R\left(dw_2-1,\frac{\lambda}{w_1}\middle|\chi\right),$$

where $l \geq 0$.

When $x = 0$, by Theorem 1, we get

$$\sum_{i=0}^{l}\binom{l}{i}\mathcal{E}_{i,\frac{\lambda}{w_1},\chi}w_1^i w_2^{l-i}R\left(dw_1-1,\frac{\lambda}{w_2}\middle|\chi\right)$$
$$= \sum_{i=0}^{l}\binom{l}{i}\mathcal{E}_{i,\frac{\lambda}{w_2},\chi}w_2^i w_1^{l-i}R\left(dw_2-1,\frac{\lambda}{w_1}\middle|\chi\right), \quad (l \geq 0).$$

By (2.5), we get

$$2I_{\chi}\left(w_1, w_2 \mid \lambda\right)$$
$$= \sum_{l=0}^{dw_2-1}\left(-1\right)^l\chi\left(l\right)\int_X\left(1+\lambda t\right)^{\frac{w_2}{\lambda}\left(w_2+w_1x+\frac{w_1}{w_2}l\right)}\chi\left(x_2\right)d\mu_{-1}\left(x\right)$$
$$= \sum_{n=0}^{\infty}\left(\sum_{l=0}^{dw_2-1}\left(-1\right)^l\chi\left(l\right)\mathcal{E}_{n,\frac{\lambda}{w_2},\chi}\left(w_1x+\frac{w_1}{w_2}l\right)w_2^n\right)\frac{t^n}{n!}. \tag{2.9}$$

On the other hand,

$$2I_{\chi}\left(w_2, w_1 \mid \lambda\right) = 2I_{\chi}\left(w_1, w_2 \mid \lambda\right)$$
$$= \sum_{n=0}^{\infty}\left(\sum_{l=0}^{dw_1-1}\left(-1\right)^l\chi\left(l\right)\mathcal{E}_{n,\frac{\lambda}{w_1},\chi}\left(w_2x+\frac{w_2}{w_1}l\right)w_1^n\right)\frac{t^n}{n!}. \tag{2.10}$$

Therefore, by (2.9) and (2.10), we obtain the following theorem.

Theorem 2. *For $w_1, w_2, d \in \mathbb{N}$ with $d \equiv 1 \pmod 2$, $w_1 \equiv 1 \pmod 2$ and $w_2 \equiv 1 \pmod 2$, let χ be a Dirichlet character with conductor d. Then, we have*

$$w_2^n\sum_{l=0}^{dw_2-1}\left(-1\right)^l\chi\left(l\right)\mathcal{E}_{n,\frac{\lambda}{w_2},\chi}\left(w_1x+\frac{w_1}{w_2}l\right)$$
$$= w_1^n\sum_{l=0}^{dw_1-1}\left(-1\right)^l\chi\left(l\right)\mathcal{E}_{n,\frac{\lambda}{w_1},\chi}\left(w_2x+\frac{w_2}{w_1}l\right), \quad (n \geq 0).$$

To derive some interesting identities of symmetry for the generalized degenerate Euler polynomials attached to χ, we used the symmetric properties for certain fermionic p-adic integrals on $\mathbb{Z}_p$. When $w_2 = 1$, from Theorem 2, we have

$$\begin{aligned}&\sum_{l=0}^{d-1} (-1)^l \chi(l) \mathcal{E}_{n,\lambda,\chi}(w_1 x + w_1 l)\\ &= w_1^n \sum_{l=0}^{dw_1-1} (-1)^l \chi(l) \mathcal{E}_{n,\frac{\lambda}{w_1},\chi}\left(x + \frac{1}{w_1} l\right).\end{aligned}$$

In particular, for $x = 0$, we get

$$\begin{aligned}&\sum_{l=0}^{d-1} (-1)^l \chi(l) \mathcal{E}_{n,\lambda,\chi}(w_1 l)\\ &= w_1^n \sum_{l=0}^{dw_1-1} (-1)^l \chi(l) \mathcal{E}_{n,\frac{\lambda}{w_1},\chi}\left(\frac{1}{w_1} l\right).\end{aligned}$$

References

1. S. Araci, A. Bagdasaryan, C. Özel, and H. M. Srivastava, *New symmetric identities involving q-zeta type functions*, Appl. Math. Inf. Sci. **8** (2014), no. 6, 2803–2808. MR 3228678
2. A. Bayad and J. Chikhi, *Apostol-Euler polynomials and asymptotics for negative binomial reciprocals*, Adv. Stud. Contemp. Math. (Kyungshang) **24** (2014), no. 1, 33–37. MR 3157406
3. L. Carlitz, *Degenerate Stirling, Bernoulli and Eulerian numbers*, Utilitas Math. **15** (1979), 51–88. MR 531621 (80i:05014)
4. R. Dangi, M. Tiwari, and C. L. Parihar, *Generalized Euler polynomials and their properties*, J. Rajasthan Acad. Phys. Sci. **12** (2013), no. 4, 385–392. MR 3299610
5. L. Jiu, V. H. Moll, and C. Vignat, *Identities for generalized Euler polynomials*, Integral Transforms Spec. Funct. **25** (2014), no. 10, 777–789. MR 3230659
6. D. S. Kim, *Identities associated with generalized twisted Euler polynomials twisted by ramified roots of unity*, Adv. Stud. Contemp. Math. (Kyungshang) **22** (2012), no. 3, 363–377. MR 2976595
7. D. S. Kim, N. Lee, J. Na, and K. H. Park, *Identities of symmetry for higher-order Euler polynomials in three variables (I)*, Adv. Stud. Contemp. Math. (Kyungshang) **22** (2012), no. 1, 51–74. MR 2931605
8. T. Kim, *Symmetry identities for the twisted generalized Euler polynomials*, Adv. Stud. Contemp. Math. (Kyungshang) **19** (2009), no. 2, 151–155. MR 2566912 (2010j:11041)
9. ———, *Symmetry of power sum polynomials and multivariate fermionic p-adic invariant integral on $\mathbb{Z}_p$*, Russ. J. Math. Phys. **16** (2009), no. 1, 93–96. MR 2486809 (2010c:11028)
10. ———, *An identity of symmetry for the generalized Euler polynomials*, J. Comput. Anal. Appl. **13** (2011), no. 7, 1292–1296. MR 2791956 (2012d:11050)
11. K. Kozuka, *On a p-adic interpolating power series of the generalized Euler numbers*, J. Math. Soc. Japan **42** (1990), no. 1, 113–125. MR 1027544 (90j:11020)

12. Q.-M. Luo, *q-analogues of some results for the Apostol-Euler polynomials*, Adv. Stud. Contemp. Math. (Kyungshang) **20** (2010), no. 1, 103–113. MR 2597996 (2011e:05031)
13. Q.-M. Luo and F. Qi, *Relationships between generalized Bernoulli numbers and polynomials and generalized Euler numbers and polynomials*, Adv. Stud. Contemp. Math. (Kyungshang) **7** (2003), no. 1, 11–18. MR 1981601
14. S.-H. Rim, J.-H. Jeong, S.-J. Lee, Eun-Jung Moon, and Joung-Hee Jin, *On the symmetric properties for the generalized twisted Genocchi polynomials*, Ars Combin. **105** (2012), 267–272. MR 2976377
15. C. S. Ryoo, *On the generalized Barnes type multiple q-Euler polynomials twisted by ramified roots of unity*, Proc. Jangjeon Math. Soc. **13** (2010), no. 2, 255–263. MR 2676690 (2011e:11043)
16. E. Şen, *Theorems on Apostol-Euler polynomials of higher order arising from Euler basis*, Adv. Stud. Contemp. Math. (Kyungshang) **23** (2013), no. 2, 337–345. MR 3088764
17. Y. Simsek, O. Yurekli, and V. Kurt, *On interpolation functions of the twisted generalized Frobenius-Euler numbers*, Adv. Stud. Contemp. Math. (Kyungshang) **15** (2007), no. 2, 187–194. MR 2356176 (2008g:11193)
18. S. L. Yang and Z. K. Qiao, *Some symmetry identities for the Euler polynomials*, J. Math. Res. Exposition **30** (2010), no. 3, 457–464. MR 2680613 (2011d:11045)
19. Z. Zhang and H. Yang, *Some closed formulas for generalized Bernoulli-Euler numbers and polynomials*, Proc. Jangjeon Math. Soc. **11** (2008), no. 2, 191–198. MR 2482602 (2010a:11036)

Department of Mathematics, Sogang University, Seoul 121-742, Republic of Korea
E-mail address: dskim@sogang.ac.kr

Department of Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea
E-mail address: tkkim@kw.ac.kr